

Single Molecule Magnetic Resonance and Quantum Computation

Haiqing Wei and Xin Xue

Department of Electrical and Computer Engineering, McGill University

Montreal, Quebec, Canada H3A 2A7

dhw@tsp.ece.mcgill.ca

xxue@photonics.ece.mcgill.ca

It is proposed that nuclear (or electron) spins in a trapped molecule would be well isolated from the environment and the state of each spin can be measured by means of mechanical detection of magnetic resonance. Therefore molecular traps make an entirely new approach possible for spin-resonance quantum computation which can be conveniently scaled up. In the context of magnetic resonance spectroscopy, a molecular trap promises the ultimate sensitivity for single spin detection and an unprecedented spectral resolution as well.

PACS numbers: 03.67.Lx, 39.90.+d, 89.80.+h

A quantum computer (QC), if ever functions, will outperform any classical computer in certain computation and simulation tasks; however building a QC which is sizeable and robust against decoherence imposes a great challenge on experimental physics, even on principle design¹. In this letter we propose that nuclear (or electron) spins in a trapped molecule would be well isolated from the environment and the state of each spin can be measured by means of mechanical detection of magnetic resonance (MD or MR). With modern MR techniques^{2,3}, a single trapped molecule amounts to an appealing piece of hardware for experimental quantum computation. In the context of MR spectroscopy, a molecular trap promises the ultimate sensitivity for single spin detection and an unprecedented spectral resolution as well.

Nuclear spin systems are good for quantum computation because they are intrinsically isolated from their environment and spin-spin coupling in a natural spin lattice enables universal quantum logic operations^{4,5}. Quantum computation based on a single molecule or nanocrystal does not encounter the problem of exponential decay in signal-to-noise ratio suffered by solution NMR⁶, provided that the state of each spin can be directly measured. Unfortunately it seems very difficult to realize the nanoprobe scheme for detection and dynamical control of single nuclear spins⁷. Indeed, scientists have been pursuing the goal of single-spin detection using various methods, including optical spectroscopy on single molecules^{8,9} and the very promising magnetic resonance force microscopy (MRFM)¹⁰⁻¹². However detecting single nuclear spins with MRFM requires high-quality cantilever and extreme magnetic field gradient which are still out of the reach of today's technology. Besides, with the sample fixed on a solid, MRFM may not be suitable for NMR quantum computation due to the decoherence problem. By contrast, a single molecule or nanoparticle

suspended in an electromagnetic or optical trap constitutes a harmonic oscillator with small inertia and ultra-sensitive "spring" as well as often negligible damping. A tiny force exerted on a single spin is readily detectable. Better isolation in a molecular trap is an added bonus to quantum computation.

The extensively used Paul and Penning traps^{13,14} are suitable for storing charged molecules or particles. Actually small molecular ions even charged micro-particles have already been stored and studied in laboratory^{13,15,16}. MR study can be conveniently incorporated into the Penning trap which uses a strong uniform magnetic field for trapping. Consider a singly charged molecule containing $N \sim 10^2$ nuclei stored in the Penning trap with a magnetic field $B \sim 5$ T, its mass $M \sim 10Nm_p$ where $m_p = 1.67 \times 10^{-27}$ kg is the proton rest mass, if with a nearly spherical shape its radius $R \sim 2.52N^{1/3}a_0$ where $4a_0 = 2.12 \times 10^{-10}$ m approximates the typical separation between adjacent nuclei, then the rotational inertia $I \sim 2MR^2/5 \sim 25.4N^{5/3}m_p a_0^2$. Using the trapping parameters for a proton in Ref.14 (Table II), the translational motion of the molecular ion will be characterized by $\nu_c \sim 76$ kHz, $\nu_z \sim 318$ kHz, which are the cyclotron and axial frequencies respectively. Here one has $\nu_c < \nu_z$, which indicates the inadequate radial confinement. Another potential, such as a Paul-type RF (radio frequency) trap, is required to confine the radial motion. One is then working on a hybrid ion trap. Assuming thermal energy ~ 4.2 K, the ion undergoes classical motion with velocity $v \sim 8.3$ m/sec, occupies an orbit with diameter $\sim 30 \mu\text{m}$ in the xy plane and oscillates along the z -axis with amplitude $A = v/2\pi\nu_z \sim 4 \mu\text{m}$. The molecular rotation is hardly quantum with the angular speed $\omega \sim 2.13 \times 10^{10} \text{ sec}^{-1}$. The molecular vibration is of quantum nature because the typical phonon frequency $\nu_p \sim C/R \sim 10^{12}$ Hz amounts to higher energy than 4.2 K, here $C \sim 10^3$ m/sec is an effective sound speed. With vanishing magnetic field inhomogeneity and negligible relativistic coupling to electric fields, nuclear spins are entirely isolated from the translational motion and other external perturbations. Due to the large mismatch between ν_p and the spin precession frequency $\nu_s \sim 2\mu_N B/h \sim 76$ MHz, being $\mu_N = 5.05 \times 10^{-27}$ J/T the nuclear magneton, lattice vibration mediated interactions would be weak because they can only take place via multi-phonon processes and the fast molecular rotation ought average out the dipole-dipole coupling by the effect of motional narrowing^{3,17}. The numerical values

are sufficient to establish the feasibility of storing heavy molecular ions in electromagnetic traps and isolating nuclear spins from the environment at the same time. If stronger confinement is applied along the radial direction by, for instance, a filament electrode exerting radial electric field, heavy ions with higher energy may be trapped and probably better at preserving spin coherence. Besides, there is the possibility of applying external force to drive molecular rotation in favor of preserving quantum coherence.

To measure a specified spin using MD of MR, one applies an inhomogeneous magnetic field and an RF field in resonance to the spin Larmor precession such that the target spin flips up and down with the frequency ν_z and exerts a resonant force on the z -axial harmonic oscillator, in the same spirit as MRFM^{10–12}. However the molecular oscillator is far more sensitive than a mechanical cantilever. The effective “spring constant” is extremely small, $k = M(2\pi\nu_z)^2 \sim 6.68 \times 10^{-12}$ N/m, and the Q value is so high, $Q > 10^9$ (see Ref.14), that damping can be just neglected. Assuming a field gradient $\partial B/\partial z \sim 100$ T/m, a typical nuclear moment experiences a force $f \sim \mu_N(\partial B/\partial z) \sim 5.05 \times 10^{-25}$ N which elongates the “spring” by $f/k \sim 7.56 \times 10^{-14}$ m. The force is tiny, but mechanical resonance will significantly amplify its effect. Depending upon 0 or π phase difference, the tiny force does work $\sim \pm 4fA$ to the oscillator within each cycle. Then after about 10^7 cycles, taking time $t \sim 30$ sec, the resonant force could increase or decrease the amplitude of the z -axial ion oscillation by $\Delta A \sim (4f/k)\nu_z t \sim 3 \mu\text{m}$ which is more than observable! However, it may impose some difficulty to flip the nuclear spin at the high rate of 318 kHz which requires a strong RF field on the order of hundred gauss. It seems more appropriate to apply an oscillating magnetic gradient than to flip the spin so quickly. Then one has to find a way to eliminate the effect of other spins. For example, one may periodically, but not so quickly, flip the target spin and simultaneously invert the magnetic field gradient such that in a complete cycle of spin flip the total contribution due to other spins vanishes while the contribution from the target spin grows steadily.

Apart from ion traps, a single molecule or nanocrystal embedded in a micron-sized particle can be trapped in vacuum by means of optical levitation using merely a laser beam^{18–20}. For a micro particle with diameter $2R \sim 1 \mu\text{m}$, the typical phonon frequency C/R is on the order of GHz still much higher than the nuclear Larmor frequency. The large rotational inertia may render the particle rotation too slow at thermal energy, however with virtually no rotational damping at vacuum the particle may be driven by some intentionally applied torque to rotate at an angular frequency up to 1 GHz^{19,20}. Again nuclear spins in such a particle are well isolated from external perturbations and the dipole-dipole coupling among them can be effectively washed out by the fast driven rotation. The force sensitivity of an optical trap is extremely high so that mechanical isolation and

optical field stabilization are necessary or a feedback system should be equipped¹⁹. Quote from Ref.19 for an ultra-sensitive example: “a particle levitated at the focus of a beam having a $70 \mu\text{m}$ focal spot diameter moves $10 \mu\text{m}$ in height for a force change or gravity change of about one part in 10^6 of the particle’s weight”. If a $0.1 \mu\text{m}^3$ sized particle with mass $M \sim 2 \times 10^{-16}$ kg being levitated, the effective “spring constant” turns to be $k \sim 2 \times 10^{-16}$ N/m and the frequency of harmonic oscillation $\nu_z \sim 0.16$ Hz. In a magnetic field gradient $\partial B/\partial z \sim 500$ T/m generated from a millimeter sized ferromagnetic particle¹¹, the force exerted on a single proton comes to $f = \mu_p(\partial B/\partial z) \sim 7 \times 10^{-24}$ N which displaces the particle by $f/k \sim 35$ nm. The displacement is readily detectable by various methods used in measuring the cantilever deflection in atomic force microscopy²¹, or one may spend some longer time to accumulate the resonant oscillation and be confident of neglecting the damping¹⁸.

The numerical estimation conclusively suggests a new type of NMR spectroscopy for single molecules stored in currently used electromagnetic or optical traps, which are subject to further improvement and optimization. This will extend NMR applications to probing physical and chemical phenomena at the molecular scale. Some questions, such as spin dynamics and relaxation in isolated molecules, may only be addressed with single molecule NMR. In particular, using molecular traps it may become reality to determine molecular structure by means of nuclear spin detection, which is of great importance¹². Although the spatial resolution may be spoiled by molecular rotation and translational motion of the target molecule, different nuclei may be distinguished according to their characteristic gyromagnetic ratios and chemical environments. Moreover, high spatial resolution can be retained by inverting positions of the molecule and the source of inhomogeneous magnetic field, that is, fixing the target molecule but suspending in the trap a micro ferromagnetic ball which generates a strong inhomogeneous field²². Of course one of the most important and exciting application of molecular traps goes to spin-resonance quantum computation, provided, however, that quantum decoherence can be really suppressed to a sufficiently low rate.

It has been established that two-bit quantum gates such as exclusive-OR along with one-bit phase shift are sufficient to carry out any unitary operation²³. In the context of NMR spectroscopy, the so-called unitary quantum gates are in effect routinely realized as coherent spin manipulations^{24,25}. If complete knowledge of spin-spin coupling and the initial spin configuration in a stored molecule are available, its spin lattice constitutes a quantum processor which is conveniently described by the model of quantum cellular automaton, where quantum state swapping between any two spin sites is a basic and important operation⁴. For the trapped molecule to be really qualified as a QC, it is further assumed that a special nucleus with its uniquely recognizable gyromagnetic

ratio is identified, such that under certain RF excitation only the special spin is in resonance and flips up and down periodically while all other spins are out of resonance thus remain intact. The special nucleus will serve as the read/write port for the computer, since its spin state can be uniquely measured using MD of MR. The source of inhomogeneous magnetic field is necessarily variable in order to coordinate coherent quantum operation and decoherent quantum measurement. During coherent operation the inhomogeneous magnetic field must be completely switched off to avoid decoherence, while switching on the magnetic inhomogeneity for quantum measurement will lead to strong interaction between nuclear spins and the macroscopic measurement device which in turn results in the intentional spin state reduction. The principle of quantum measurement with MD of MR is based on a variant Stern-Gerlach effect²⁶: in the presence of the RF and inhomogeneous magnetic fields, the wave packet of a spin particle splits according to spin up or down in the reference frame rotating synchronously with the RF field, and the harmonic oscillator of the molecular trap experiences positive or negative driving force correspondingly. Depending upon the initial phase (0 or π) of the oscillator, the up or down spin state will be entangled with an enhanced or suppressed harmonic oscillation. Actually, in the inhomogeneous magnetic field, each spin exerts a static force except the read/write spin which contributes an oscillating force on/to the harmonic oscillator, due to the selective RF excitation. Therefore different spin configurations are strongly entangled with different equilibrium positions of the harmonic oscillator, the presence of magnetic inhomogeneity immediately projects the QC onto a classical state. Once that happens, the classical spin configuration stays stable, each spin adiabatically follows its local field without longitudinal relaxation¹⁷ despite perturbations due to translational motion and molecular rotation, except for the case of resonant perturbation which should be carefully avoided. Given access to the read/write spin, any other spin can be read and preset by quantum state swapping^{4,5}. Measuring the whole spin lattice effectively initializes the QC, which thereafter undergoes unitary evolution with the inhomogeneous magnetic field completely switched off. Upon completion of unitary transformation, switching on the magnetic inhomogeneity leads to wave reduction of the QC and the classical state is readily observed using MD of MR. Obviously, the NMR QC of molecular trap does not share the problem of exponential decay in signal-to-noise ratio while enjoys the great immunity to decoherence with solution NMR computation^{6,25}. There seems to be neither principle nor technical limitation in scaling up NMR quantum computation using molecular traps, if decoherence is indeed not a problem.

Finally it is worth mentioning that molecular traps are suitable to study single electron spins as well, and magnetic force detection should perform better there simply because the magnetic moment gets 10^3 times

larger. Moreover, if the electron spin system in a trapped molecule or nanoparticle can also be made robust against decoherence, then a QC based on electron spin resonance will operate much faster than its NMR counterpart.

-
- [1] For a comprehensive reference on quantum computation, see DiVincenzo, D. *et al.* (eds.) *Quantum coherence and decoherence. Proc. Roy. Soc. Lond. A* **454**, 257 - 486 (1998).
 - [2] Ernst, R. R., Bodenhausen, G. & Wokaun, A. *Principles of Nuclear Magnetic Resonance in One and Two Dimensions* (Oxford University Press, 1987).
 - [3] Slichter, C. P. *Principles of Magnetic Resonance*, 3rd ed. (Springer-Verlag, 1992).
 - [4] Lloyd, S. A potentially realizable quantum computer. *Science* **261**, 1569 (1993).
 - [5] Wei, H., Xue, X. & Morgera, S. D. NMR quantum automata in doped crystals. E-print quant-ph/9805059 at <http://xxx.lanl.gov/>.
 - [6] Warren, W. S. The usefulness of NMR quantum computing. *Science* **277**, 1688 (1997).
 - [7] DiVincenzo, D. Quantum computation. *Science* **270**, 257 (1995).
 - [8] Wrachtrup, J. *et al.* Detection of a single electron spin. In Marti, O. & Möller, R. (eds.) *Photons and Local Probes* (Kluwer Academic Publishers, 1995).
 - [9] Gruber, A. *et al.* Scanning confocal optical microscopy and magnetic resonance on single defect centers. *Science* **276**, 2012 (1997).
 - [10] Rugar, D., Yannoni, C. S. & Sidles, J. A. Mechanical detection of magnetic resonance. *Nature* **360**, 563 (1992).
 - [11] Rugar, D. *et al.* Force detection of nuclear magnetic resonance. *Science* **264**, 1560 (1994).
 - [12] Sidles, J. A. *et al.* Magnetic resonance force microscopy. *Rev. Mod. Phys.* **67**, 249 (1995).
 - [13] Dehmelt, H. G. Radiofrequency spectroscopy of stored ions. *Advances in Atomic and Molecular Physics* **3**, 53 (1967), and **5**, 109 (1969).
 - [14] Brown, L. S. & Gabrielse, G. Geonium theory: Physics of a single electron or ion in a Penning trap. *Rev. Mod. Phys.* **58**, 233 (1986).
 - [15] DiFilippo, F. *et al.* Accurate atomic masses for fundamental metrology. *Phys. Rev. Lett.* **73**, 1481 (1994). Also see the *RLE Progress Report* **140** (MIT, 1998). Page 217 - 220.
 - [16] Wuerker, R. F., Shelton, H. & Langmuir, R. V. Electrodynamic containment of charged particles. *J. Appl. Phys.* **30**, 342 (1959).
 - [17] Abragam, A. *The Principles of Nuclear Magnetism* (Oxford University Press, 1961).
 - [18] Ashkin, A. & Dziedzic, J. M. Optical levitation in high vacuum. *Appl. Phys. Lett.* **28**, 333 (1976).
 - [19] Ashkin, A. & Dziedzic, J. M. Feedback stabilization of optically levitated particles. *Appl. Phys. Lett.* **30**, 202 (1977).
 - [20] Ashkin, A. Applications of laser radiation pressure. *Science* **210**, 1081 (1980).
 - [21] For example, Rugar, D., Mamin, H. J. & Guethner,

- P. Improved fiber-optic interferometer for atomic force microscopy. *Appl. Phys. Lett.* **55**, 2588 (1989).
- [22] Xue, X. & Wei, H. (in preparation).
- [23] Barenco, A. *et al.* Elementary gates for quantum computation. *Phys. Rev. A* **52**, 3457 (1995).
- [24] Cory, D. G., Price, M. D. & Havel, T. F. Nuclear magnetic resonance spectroscopy: an experimentally accessible paradigm for quantum computing. *Physica D*, to appear. E-print quant-ph/9709001 at <http://xxx.lanl.gov/>.
- [25] Gershenfeld, N. A. & Chuang, I. L. Bulk spin-resonance quantum computation. *Science* **275**, 350 (1997).
- [26] Sidles, J. A. Folded Stern-Gerlach experiment as a means for detecting nuclear magnetic resonance in individual nuclei. *Phys. Rev. Lett.* **68**, 1124 (1992).